

CANNONADE 212 EC

HERBICIDE

212g/L FLUAZIFOP-P



Defend your
broadleaf crops



for superior
grassy weed
control in
broadleaf crops

SureFire

CANNADE 212 EC

HERBICIDE

212g/L FLUAZIFOP-P

SureFire CANNADE 212EC

herbicide is a selective phenoxy herbicide used for post-emergence control of annual and perennial grass weeds, but does little or no harm to broad-leaved plants. It is rapidly absorbed by leaves and green stems and translocated throughout the plant.

It accumulates at growing points, both above ground and in the roots, rhizomes and stolons of grass weeds. Weed growth, and hence competition with the crop, stops within 48 hours.

It has an extensive label for use on soybeans and other broad-leaved crops such as carrots, spinach, potatoes, and ornamentals, fruit orchards & pine plantations.

SureFire CANNADE212EC features:

1. Selective and systemic post-emergent herbicide.
2. Controls many annual and perennial weeds as shown typically below (refer to the label for specifics).
3. Non-toxic to broadleaf crops.
4. Resistant to wash-off by rain, due to systemic activity.
5. Safe to birds, fish, bees, and other beneficial insects.
6. Registered for aerial applications.
7. Non-Ionic Organic Surfactant use is recommended.

CROP/ TYPICAL SITUATION	TYPICAL WEEDS CONTROLLED	WEED STAGE	RATE /HA
Canola, chickpeas, lucerne, forage & seed crops, blueberries, soybeans, hops, blackcurrants, peas, peanuts, broccoli, cabbage, cauliflower, tomatoes, lettuce, celery, carrots, sunflowers, apples, pears, grapes, stonefruit, citrus, mangoes, avocados, bananas, pawpaw, guava, onions, potatoes, cucurbits, strawberries, pineapples, potted & ornamental shrubs, pine plantations, oilseed poppies, duboisia	Barnyard Grass, Crowsfoot Grass, Liverseed Grass, Summer Grass (Crabgrass), Stinkgrass, Guinea Grass, Annual (Wimmera) Ryegrass, Wild Oats, Barley Grass, Brome Grasses, Annual Phalaris, Volunteer Cereals, seedlings of Couch Grass, English Couch (rope twitch), and Water Couch, Johnson Grass (seedling) and Paspalum (seedling)	Generally growing actively at early growth stages	Typically 500mL to 4L depending on grass to be controlled (higher rates for couch, paspalum, kikuyu). <i>..always refer to product label for specific mixing & application rates.</i>
	Established plants of Couch Grass, English Couch (rope twitch), Water Couch, Johnson Grass, Paspalum, Bent Grass, Kikuyu Grass and Kentucky Bluegrass		



Container size:	1 L	Carton quantity:	5
	5 L		4
	20 L		

GROUP A HERBICIDE APVMA approval No 85320/112899

© PCT International Pty Ltd

www.pctrural.com.au