

# MACROMITTE

**MITICIDE**

480 g/L BIFENAZATE

**NOW WITH  
ALMONDS ,  
STRAWBERRIES  
& FRUITING  
VEGETABLES  
ON THE LABEL !**



## **SUPERIOR MITE CONTROL**

- Unique chemistry & mode of action.
- Ideal for IPM.
- Additional label claims.

**SûreFire**

# MACROMITE

## MITICIDE

SureFire MACROMITE Miticide is a selective miticide for the control of plant-eating mite pests on apples, pears, apricots, nectarines, peaches and plums, cucumber, fruiting vegetables, papaya, peppers, zucchini, eggplant, bitter melon, sin qua, tomatoes, snake beans, hops, and numerous varieties of berries\*\*.

When used as directed and applied to the foliage, it provides quick knockdown through contact activity, and good residual control.

Due to its unique chemistry, Group 20D mode of action and selective nature, SureFire Macromite Miticide is relatively inactive against beneficial and/or predatory mites and insects and therefore is compatible with IPM and resistance management programs.

\*\*as per APVMA Permit Nos : PER82341, PER14425 & PER13265. Refer to these permits for specific application rates, withholding periods & constraints.

Crop	Pest	Rate	WHP	Critical Comments
Apples & Pears	Two-spotted mite ( <i>Tetranychus urticae</i> ) European red mite	Dilute spraying 65 mL/100 L	Harvest 7 days	Surefire Macromite Miticide should be applied to the point of run-off as soon as mites appear. Equipment should be calibrated so that no less than 1000 L of water is applied per hectare. Complete coverage of all foliage is essential. Only one application should be made per season.
Apricots, nectarines, peaches & plums	( <i>Panonychus ulmi</i> ) Bryobia mite ( <i>Bryobia rubrioculus</i> )		Grazing 28 days	
			Harvest 3 days	
Almonds	Two-spotted mite ( <i>Tetranychus urticae</i> ) Bryobia mite ( <i>Bryobia rubrioculus</i> )		Grazing 28 days	
Fruiting vegetables including; tomatoes, capsicum, peppers and eggplant (excluding sweet corn and mushrooms)			Harvest 14 days	
			Grazing 28 days	
			1 day	
Cucurbits (excluding cucumber)			3 days	
Cucumber			1 day	
Papaya			7 days	
Strawberries	Two-spotted mite ( <i>Tetranychus urticae</i> ), Bryobia mite ( <i>Bryobia rubrioculus</i> )		1 day	Do not apply more than 2 applications per growing season. Do not re-apply within 21 days of the previous application. A different acaricide should be used between Surefire Macromite applications.

Container sizes: Carton quantity:

1 litre 5

5 litre 4

Note: Only one application of SureFire Macromite should be applied per season for effective insect resistance management.  
WHP vary between 3-14 days for harvest.  
Refer to the full label for specific information.

**GROUP 20D INSECTICIDE**

APVMA Approval No: 81158/131207



© **PCT International Pty Ltd**

[www.pctrural.com.au](http://www.pctrural.com.au)