



Company Name: PCT HOLDINGS PTY LTD
Product Name: Surefire Pro Cockroach Gel Bait

APVMA Approval No: 81036/102199

Label Name:	Surefire Pro Cockroach Gel Bait
--------------------	---------------------------------

Signal Headings:	READ SAFETY DIRECTIONS BEFORE OPENING OR USING
-------------------------	--

Constituent Statements:	ACTIVE CONSTITUENT: 0.5 g/kg FIPRONIL
--------------------------------	---------------------------------------

Mode of Action:	<table border="1"><tr><td>GROUP</td><td>2B</td><td>INSECTICIDE</td></tr></table>	GROUP	2B	INSECTICIDE
GROUP	2B	INSECTICIDE		

Statement of Claims:	For the treatment of cockroach infestations in commercial, industrial and residential premises.
-----------------------------	---

Net Contents:	15 g 30 g 60 g 120 g (4 x 30 g)
----------------------	--

Restraints:	RESTRAINTS: DO NOT place in areas that are routinely washed. DO NOT apply this product in any areas easily accessible to children or animals.
--------------------	---

Directions for Use:	This section contains file attachment. File Name: Surefire Pro Gel label DfU 0315.doc File Size: 32768 bytes
----------------------------	--

Other Limitations:	NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.
---------------------------	---

Withholding Periods:	
-----------------------------	--

Trade Advice:	
General Instructions:	<p>GENERAL INSTRUCTIONS</p> <p>To apply the gel, remove the cap on the nozzle and touch the tip to the surface to be treated and slowly depress plunger until sufficient gel is dispensed. Recap the dispenser after treatment is completed.</p> <p>Where maintenance treatments form part of the professional pest controllers contract it is recommended that cockroach activity be monitored where gel has been applied one month after initial treatment or at more regular intervals in areas of high infestation.</p>
Resistance Warning:	<p>Insecticide Resistance Warning GROUP 2B INSECTICIDE</p> <p>For insecticide resistance management SUREFIRE PRO Cockroach Gel Bait is a Group 2B Insecticide. Some naturally occurring insect biotypes resistant to SUREFIRE PRO Cockroach Gel Bait and other group 2B insecticide may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if this product or other group 2B insecticides are used repeatedly. The effectiveness of this product on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, PCT Holdings Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant insects. This product may be subject to specific resistance management strategies. For further information, contact your local supplier, PCT Holdings Pty Ltd representatives or local agricultural department agronomist.</p>
Precautions:	
Protections:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p>
Storage and Disposal:	<p>STORAGE AND DISPOSAL</p> <p>Store in the closed, original container in a cool, well ventilated locked place out of the reach of children. Do not store in direct sunlight and do not store at temperatures exceeding 35°C. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in the garbage.</p>
Safety Directions:	<p>SAFETY DIRECTIONS</p> <p>May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.</p>
First Aid Instructions:	<p>FIRST AID</p> <p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.</p>
First Aid Warnings:	

SITUATION	PEST	RATE
Domestic, commercial and public service buildings including: farm buildings, food processing establishments, factories, hospitals, hospitals, homes, kitchens, offices, restaurants, retail outlets, storerooms, vehicles and other situations which fall within these definitions.	German and other common cockroaches.	<p>Apply at least 2 spots per m². Increasing the number of spots (up to 5/m²) may be required for heavy infestations or if the American cockroach is the main pest.</p> <p>Recommended spot size is approximately 5 mm diameter (0.1 g) or the size of the spot indicated below:</p> <p style="text-align: center;">●</p>
CRITICAL COMMENTS		
<p>Conduct a thorough inspection of the premises and then place gel in areas near cockroach harbourages or in sheltered areas where signs of cockroaches are evident.</p> <p>Replenish gel if previous application has been consumed or cockroaches remain a problem; increasing number of spots if necessary.</p>		