

AZLAN

FUNGICIDE

ACTIVE CONSTITUENTS: 200 g/L TEBUCONAZOLE
120 g/L AZOXYSTROBIN



**for foliar diseases on
grapevines**

SureFire

AZLAN

FUNGICIDE

**ACTIVE CONSTITUENTS: 200 g/L TEBUCONAZOLE
120 g/L AZOXYSTROBIN**

SUREFIRE AZLAN is a very effective product for control of powdery mildew and downy mildew in grapevines, suitable for either dilute spraying or concentrate spraying.

It can be applied as part of a season-long spray program, targeting all growth stages from shoots to after fruit set.

However, it should not be applied at more than 2 applications per season, to avoid fungicide resistance issues.

Withholding period for this product is 4 weeks after application to harvest.

Refer label for specific information.

Off-label Permits applicable for AZLAN

Permit Number	Expires	Crops	Pest
PER82104	Nov 22	Adzuki, Mung, Navy Beans	Powdery Mildew
PER89356	Sep 23	Maize	Northern Corn Leaf Blight
PER89395	May 22	Soya Beans	Anthraxnose
PER86245	Feb 26	Sweet Corn	Maize Rust
PER13717	May 22	Tomatoes	Powdery Mildew

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Grapevines	Powdery Mildew (<i>Erysiphe necator</i>) (formerly <i>Uncinula necator</i>)	Dilute Spraying: 65 or 100 mL/100 L Concentrate spraying: 1 L/ha	H 4 weeks	Apply Surefire Aztec Fungicide thoroughly as part of a season-long spray program targeting the following growth stages: 1. When shoots are 20 cm long. 2. Pre-flowering. 3. Flowering. 4. After fruit set. 5. Before bunches close. Use the lower rate prior to flowering (before E-L 19) or under low disease pressure. Apply the higher rate from the start of flowering (from E-L 19) and under conditions of high disease pressure. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. DO NOT apply in volumes less than 250 L/ha. DO NOT apply at concentrations greater than 4X maximum label rate. This use is subject to a CropLife Fungicide Resistance Management Strategy: DO NOT apply more than two sprays of Surefire Aztec Fungicide per season. DO NOT use Surefire Aztec Fungicide curatively.
	Downy Mildew (<i>Plasmopara viticola</i>)	Dilute spraying: 100 mL/ 100 L Concentrate spraying: 1 L/ha		Apply Surefire Aztec Fungicide as part of a seasonal preventative spray program commencing when shoots are approximately 20 cm long. Use Surefire Aztec Fungicide in rotation with other mode of action fungicides on a 7-21 day spray interval. Under conditions of high disease pressure and/or at critical timings i.e. during flowering; ensure that Surefire Aztec Fungicide is applied in rotation with fungicides from a different mode of action group.
	Botrytis Bunch Rot (<i>Botrytis cinerea</i>)			When Surefire Aztec Fungicide is used in a seasonal spray programme it will provide control of Botrytis additional to that of specific botryticides. Under conditions of high disease pressure, Surefire Aztec Fungicide should not be used for Botrytis control particularly at critical timings such as 80 -100% capfall and pre-harvest. This use is subject to a CropLife Fungicide Resistance Management Strategy: Apply Surefire Aztec Fungicide preventatively. DO NOT apply more than two sprays per season of Surefire Aztec Fungicide or any other Group 11 fungicide. If three or fewer Botrytis sprays are applied in a season, use no more than one spray of Surefire Aztec Fungicide during the season. If four or more Botrytis sprays are to be applied in a season, up to two Surefire Aztec Fungicide sprays can be applied per season.

H4 harvest

Container sizes: Carton quantity:

1 litre 5

5 litre 4

Note: This product is phytotoxic to certain apple varieties — avoid spray drift near apple trees & do not use spray equipment on apple trees. Refer to the label for specific information.

GROUP 3 11 FUNGICIDE

APVMA Approval No: 87767/131989



© PCT International Pty Ltd

www.pctrural.com.au