

EXCLAIM 44 SG

INSECTICIDE

44 g/Kg EMAMECTIN

for control of:

diamondback moth

white butterfly

heliothis

in selected crops



SûreFire

SureFire EXCLAIM 44 SG

INSECTICIDE

44 g/Kg EMAMECTIN

SureFire EXCLAIM 44 SG Insecticide is our new emamectin based insecticide registered for control of diamondback moth, cabbage moth, cabbage white butterfly, and heliothis (see label below).

It is supplied as a soluble granular formulation, featuring translaminar activity, rapid rainfastness and minimal effect on beneficial insects.

CROP	PEST	RATE	CRITICAL COMMENTS
Brassica Vegetables Broccoli, Brussels Sprouts, Cabbages, Cauliflower only	Diamondback Moth (Cabbage Moth) (<i>Plutella xylostella</i>) and Cabbage White Butterfly (<i>Pieris rapae</i>)	250 to 300 g/ha	Spray at first signs of insect infestation as indicated by local spray thresholds. Use the lower rate on low to moderate infestations. Add Citowett* or Agral® at the appropriate label rate. DO NOT make more than 4 applications to any brassica crop. Where more than one crop is grown DO NOT make more than 4 applications of Surefire Exclaim in any one year. Surefire Exclaim should be used according to CropLife insecticide resistance strategies.
Capsicum, Lettuce	Heliothis (<i>Helicoverpa armigera</i> and <i>Helicoverpa punctigera</i>)	150 to 250 g/ha	Spray at first signs of insect infestation as indicated by local spray thresholds. For best results apply soon after the <i>Helicoverpa</i> eggs have hatched. DO NOT apply more than 4 sprays of Surefire Exclaim per crop. Where more than one crop is grown per year DO NOT apply more than 4 sprays per year. Ensure thorough spray coverage. Use the lower rate when targeting light infestations of small larvae. Use 250 g/ha during periods of heavy insect pressure or under very hot and sunny conditions. Surefire Exclaim should be used according to the CropLife resistance strategies.
Tomatoes	Heliothis (<i>Helicoverpa armigera</i> , <i>Helicoverpa punctigera</i>)	150 to 250 g/ha or 15 to 25 g/100 L (dilute spraying) for trellised Tomatoes. Concentrate Spraying Refer to the Mixing/ Application section	Spray at first signs of insect infestation as indicated by local spray thresholds. For best results apply soon after the <i>Helicoverpa</i> eggs have hatched. DO NOT apply more than 4 sprays of Surefire Exclaim per crop. Where more than one crop is grown per year DO NOT apply more than 4 sprays per year. Ensure thorough spray coverage. Use the lower rate when targeting light infestations of small larvae. Use 250 g/ha during periods of heavy insect pressure or under very hot and sunny conditions. Trellised Tomatoes: In the case of dilute spraying (g/100 L) apply in the range of 400 to 500 L/ha after transplanting and increase to 800 to 1000 L/ha at full canopy. Surefire Exclaim should be used according to the CropLife resistance strategies.

Container sizes:
500g 2.5Kg

Carton quantity:
6 4

Withholding Periods:

WITHHOLDING PERIODS

Brassica Vegetables: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION
DO NOT USE TREATED CROP, CROP WASTE OR PRODUCE FOR STOCK FOOD
Capsicum, Lettuce, Tomatoes: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION
DO NOT ALLOW LIVESTOCK TO GRAZE TREATED CROPS OR CROP STUBBLE



APVMA Approval No: 86238/115093

GROUP 6 INSECTICIDE

© **PCT International
Pty Ltd**

www.pctrural.com.au