ZINAM 500SC

FUNGICIDE

500 g/L FLUAZINAM



club root brassicas white rootapples leaf blight grapevine late blight potatoes



SuréFire

ZINAM 500SC

FUNGICIDE

500 g/L FLUAZINAM

Being a Group 29 fungicide, Zinam becomes an effective addition to your fungicide rotation strategies. It assists with your program of resistance management.

Zinam fungicide inhibits the formation and movement of spores, thereby stopping disease before it takes over.

Zinam has a multi-site mode of action that disrupts energy production in fungi. Zinam has great protective activity, possesses very good residual effect and has rain fastness.

Zinam can protect brassica crops from the spread of clubroot when used as a seedling drench in the soil.

Crop	Disease	Rate
Apples (Qld, NSW, WA & Tas only)	White Root Rot (Rosellinia necatrix)	50 mL/ 100 L
Broccoli Brussels sprouts Cabbage Cauliflower Kohlrabi	Clubroot (Plasmodiophora brassicae)	Seedling drench 25-50 mL per 1000 plants
Grape vines	Phomopsis Cane and Leaf Blight (Phomopsis viticola)	Application 100 mL / 100 L
Potatoes	Late blight (Phytophthora infestans) Sclerotinia (Sclerotinia spp.)	250 mL/ha

APVMA Approval №: 85676 / 130770

GROUP 29 FUNGICIDE

- Low use rates with unique Mode of Action
- Harvest withholding period is not required when product is used as per the label.
- Registered for use either as a seedling drench or as a pre-transplant soil application, for maximum flexibility
- ◆Treated crops must not be grazed or fed to animals.
- ♦ Adjuvants, such as Wetter 600 at 20mL/100L, or summer oil at 2% will assist coverage and contact.



Container Carton sizes: quantity:

5 litre 4

© PCT International
Pty Ltd

www.pctrural.com.au